Helmholtz-Zentrum Dresden-Rossendorf (HZDR)



Cerebral blood flow of the frontal lobe in untreated children with trigonocephaly vs healthy controls: an arterial spin labeling study

de Planque, C. A.; Petr, J.; Gaillard, L.; Mutsaerts, H. J.; van Veelen, M. L. C.; Versnel, S. L.; Dremmen, M.; Mathijssen, I. M. J.;

Originally published:

January 2022

Plastic and Reconstructive Surgery 149(2022)4, 931-937

DOI: https://doi.org/10.1097/PRS.00000000008931

Perma-Link to Publication Repository of HZDR:

https://www.hzdr.de/publications/Publ-31681

Release of the secondary publication on the basis of the German Copyright Law § 38 Section 4.

CC BY-NC

- 1 Cerebral blood flow of the frontal lobe in untreated children with trigonocephaly vs healthy controls:
- 2 an arterial spin labeling study
- 3 C.A. de Planque, MD¹ J. Petr, PhD², L. Gaillard, MD¹, H.J. Mutsaerts, MD PhD³, M.L.C. van Veelen, MD
- 4 PhD⁴S.L. Versnel, MD PhD,¹ M. Dremmen, MD, ⁵, I.M.J. Mathijssen MD, PhD, MBA-H¹
- ⁵ ¹ Erasmus MC-Sophia Children's Hospital, University Medical Center Rotterdam, Department of Plastic,
- 6 *Reconstructive Surgery and Hand Surgery, Rotterdam, The Netherlands.*
- 7 ² Helmholtz-Zentrum Dresden-Rossendorf, Institute of Radiopharmaceutical Cancer Research, Germany
- 8 ³Amsterdam UMC/UMC Utrecht, The Netherlands.
- 9 ⁴ Erasmus MC-Sophia Children's Hospital, University Medical Center Rotterdam, Department of
- 10 *Neurosurgery, Rotterdam, The Netherlands.*
- ⁵ Erasmus MC-Sophia Children's Hospital, University Medical Center Rotterdam, Department of
- 12 Radiology and Nuclear Medicine, Rotterdam, The Netherlands

- 14 **Corresponding Author:**
- 15 C.A. de Planque, MD
- 16 Email: <u>c.deplanque@erasmusmc.nl</u>
- 17 Adress: Ee 1591b
- 18 Dr. Molewaterplein 40
- 19 3015 GD Rotterdam, The Netherlands
- 20 Phone: +311070443291
- 21

- 22 Key words: ASL, trigonocephaly, brain, perfusion, CBF, craniosynostosis
- 23 **Financial disclosure statement**: Research of C.A.P. was supported by Sophia Stichting Wetenschappelijk
- 24 Onderzoek (project number: B-16-03a); they had no involvement in any aspect of the study.

26 Abstract

27 Background

- 28 Craniofacial surgery is the standard of treatment for children with moderate to severe trigonocephaly.
- 29 However, assessing the risk of suboptimal neurodevelopment and added value of surgery is difficult in
- 30 individual cases. In this study we aim to address the hypothesis that brain development is restricted in
- 31 trigonocephaly patients by investigating cerebral blood flow in the frontal lobe.

32 Methods

- 33 Between 2018 and 2020, we prospectively included trigonocephaly patients for whom a surgical
- 34 correction was considered in an MRI study measuring cerebral perfusion with arterial spin labeling (ASL).
- 35 The mean value of cerebral blood flow (CBF) in the frontal lobe was calculated for each subject and

36 compared between the trigonocephaly patients and healthy controls.

37 Results

- 38 MRI scans of 36 trigonocephaly patients (median age 0.5y, IQR 0.3, 11 females) were included and
- 39 compared with 16 controls without cerebral pathology (median age 0.83y, IQR 0.56, 10 females). The
- 40 mean CBF values in the frontal lobe of the trigonocephaly patients (73.0 ml/100g/min) did not appear to
- 41 be significantly different in comparison with controls (70.5 ml/100g/min, p = 0.6479). The superior,
- 42 middle, and inferior part of the frontal lobe showed no significant differences either.

43 Conclusions

- 44 Before surgery, the frontal lobe of trigonocephaly patients aged under 18 months old has a normal CBF.
- 45 In addition to the previously reported very low prevalence of papilledema or impaired skull growth, this

- 46 finding further supports our hypothesis that craniofacial surgery for trigonocephaly is rarely indicated
- 47 for signs of raised intracranial pressure.

- 49 Key words: trigonocephaly, craniosynostosis, brain perfusion, cerebral blood flow, , frontal lobe, arterial
- 50 spin labeling, ASL
- 51

52 Introduction

53 Trigonocephaly is the second most common form of non-syndromic craniosynostosis, with prenatal 54 closure of the metopic suture.¹ Patients present within a sliding scale of severity in phenotype, 55 depending on the timing of suture closure. It still remains a subject of discussion, which degree of 56 severity is clinically relevant for a surgical indication. Nowadays, patients with moderate and severe 57 phenotypes undergo surgical correction of the frontal bones and supra orbital rims aiming for 58 unrestricted brain development, to reduce the risk of neurodevelopmental disorders, and improved 59 esthetics. However, preoperatively less than 2% of the trigonocephaly patients have papilledema as a 60 sign of intracranial hypertension.² Moreover, preoperative trigonocephaly patients show a completely 61 normal intracranial volume in comparison to healthy aged-matched controls.³ Lastly, it was shown that patients with trigonocephaly are at risk of developing mental deficiencies/disorders, behavioral 62 63 problems, delays in speech and language, irrespective of having undergone surgery.^{4, 5} Taking all the 64 above together, it remains unknown what the added value of surgery is in trigonocephaly with respect 65 to future brain development.

66 The exact mechanism of the association between trigonocephaly and suboptimal neurodevelopmental 67 outcome is not fully understood.⁶ Although some have suggested that brain development is impaired as 68 a result of the synostosis, others hypothesize the increased prevalence of neurodevelopmental 69 disorders in these patients is caused by an intrinsic brain disorder. The former hypothesis of brain development restriction due to synostosis could be reflected in altered cerebral blood flow (CBF). 7-10 70 71 Brain perfusion in trigonocephaly patients was examined previously with single-photon emission 72 computed tomography (SPECT) and a lower perfusion was reported in the frontal lobe preoperatively compared to postoperatively and to the rest of the brain.^{9, 10} These two studies offered, however, 73 74 qualitative evaluation of relative perfusion values only. Over the last decade, more advanced imaging

techniques have been developed to measure cerebral perfusion, hence, this original claim was not
reassessed yet. Arterial spin labeling (ASL) Magnetic Resonance Imaging (MRI) is a technique that
provides injection-free measurements of absolute brain perfusion with quantitative accuracy
comparable to that of PET.¹¹ ASL has previously been used for several pediatric applications, but
craniosynostosis patients might present a further challenge given the skull deformations of these
patients.^{8 12, 13}

81 The aim of this study is to reassess the previous claims of perfusion changes in craniosynostosis subjects 82 to gain more insight into the hypothesis of brain restriction by investigating cerebral blood flow by 83 acquiring ASL MRI brain scans of young children with trigonocephaly preoperatively and of aged-84 matched healthy controls. Brain perfusion will be assessed in several brain regions focusing mostly, due 85 to the shape of the skull in trigonocephaly patients as well as the increased prevalence of behavior and 86 cognitive disorders, on the frontal lobe. Based on the very low prevalence of papilledema or impaired 87 skull growth in trigonocephaly compared to other craniosynostosis patients, we hypothesize that there are no abnormalities in frontal lobe perfusion in the first two years of life.^{2,3} 88

90 Methods

91 The Ethics Committee of the Erasmus Medical Center approved this prospective imaging study in patients
92 with trigonocephaly (METC-2018-124), which is part of an ongoing work at the Dutch Craniofacial Center.

93

94 Study population

95 MRI scans from children with metopic synostosis for whom a surgical correction was considered were 96 included over a period of two years (2018-2020). Surgery is only considered for moderate and severe 97 presentation, mainly defined by the forehead shape in a bird eye's view and considered present if the 98 lateral orbital rim is visible and the midline ridge is significantly prominent. This is illustrated by Birgfeld 99 et al (2013) in Figure 5.¹⁴ Children were below 2 years of age at the time of the MRI brain study. The 100 control group consisted of subjects undergoing MRI for clinical reasons. These subjects were included 101 when the following conditions were met: 1) the subjects were found to have no neurological pathology 102 of the head and neck area on imaging; 2) the subjects were free of any neurological or psychological 103 morbidity on follow-up; and 3) the subjects' MRI data were of sufficient quality to be used for research. 104

105 MRI Acquisition

All brain MRI data were acquired with a 1.5T scanner (GE Healthcare), including pseudocontinuous ASL sequences with the following imaging parameters: 3D FSE spiral readout with a stack of 8 spirals and 3 averages, TR 4604 ms, TE 10.7 ms, voxel size 3.75 x 3.75 x 4.0 mm³, axial field of view 24.0 cm, number of slices, labeling duration 1450 ms, post-labeling delay (PLD) 1025 ms, background suppression, and MO-scan acquisition for calibration. This protocol was identical in both trigonocephaly patients and controls. Both groups underwent deep sedation or anesthesia during the MRI procedure, which included using sevoflurane or propofol.

114 **ASL Data Analysis**

115 Data processing and evaluation was performed with the ExploreASL pipeline and it included the basic processing of ASL and M0 calibration images as described in the ExploreASL review paper¹⁵. T1-weighted 116 117 images were excluded from the complete analysis due to insufficient differentiation between white and 118 grey matter. This is common for the age group we studied due to incomplete myelination and it 119 precluded successful segmentation and spatial normalization in part of the subjects. To be able to use 120 the structural atlas of brain regions for the evaluation, the ASL images were directly aligned with the 121 Montreal Neurological Institute (MNI) template. In this age group, the brain is of considerably different 122 shape and size than the adult brain. Therefore, a dedicated template - UNC 0-1-2 Infant Atlases, the version for 1y - was used to replace the adult template.¹⁶ The control images were not of high-enough 123 124 contrast to allow alignment with the mean MNI T1-weighted image, therefore a registration of the 125 individual CBF to a standard-space pseudoCBF, based on the GM and WM maps of the template, was 126 performed.¹⁷ Because of the relatively high amount of skull deformations in the patient group, rigid or 127 affine registration was not considered sufficient for alignment of the individual data to the template. 128 Therefore, an affine registration followed by nonlinear deformations using a linear combination of three 129 dimensional discrete cosine transform (DCT) basis functions was used.¹⁸ The CBF has been quantified according to the consensus paper by Alsop et al.¹⁶ Whole brain and regional 130 131 CBF were evaluated in both hemispheres taken together and superior, middle, and inferior levels of the 132 frontal lobe. There were large shape and size differences in the population due to age and disease. 133 Therefore, we chose to evaluate the mean CBF values in the native space of each subject to avoid effects 134 of interpolation and voxel size change. The anatomical ROIs were taken from Hammer's atlas in the MNI 135 space and transformed to the subject's space using the previously obtained transformation for the spatial normalization.¹⁹ Mean CBF values were investigated as well for the following brain regions: 136

Frontal lobe, Occipital lobe, Parietal lobe, Temporal lobe, Insula, Cerebellum, Caudate, Putamen, andThalamus.

139

140 Statistical Analyses

141 Statistical analyses were performed using R Version 1.1.442 Parametric statistics were used when the

142 distribution of the data did not violate assumptions of normality.²⁰ Linear mixed models were used to

143 compare the mean CBF in the frontal lobe in trigonocephaly patients vs controls. To confirm the validity

of our data, we normalized CBF data of the frontal lobe by occipital lobe data, using the ratio of frontal

145 lobe/occipital lobe. We compared this ratio of frontal lobe/occipital lobe between patients and controls

by using a T-test. In addition, we assessed the ratio of the spatial coefficient of variation, a measure of

147 global spatial signal distribution, between the frontal lobe and occipital lobe, comparing the ratios

148 between patients and controls using a T-test.

149

151 Results

152

153 Patient characteristics

154 Thirty-six patients with trigonocephaly with a median age of 0.50 years (IQR 0.30) and sixteen control

subjects with a median age of 0.83 years (IQR 0.56) were included in this study as presented in Table 1.

156

157 Frontal lobe perfusion

158 By mixed model, Table 2 demonstrates that there is no significant difference in the mean perfusion of

the frontal lobe between trigonocephaly patients (73 ml/100 g/min) and controls (70.5 ml/100 g /min, p

160 = 0.6479). The mean perfusion was compared between the two groups. **Table 3** shows the mean

161 perfusion for three levels of the frontal lobe - the superior, middle, and inferior level. None of these

three levels was significantly different between the patient and control groups.

163 To account for possible variation in labelling efficiency, we chose the occipital lobe as a reference region

164 for normalization. We compared the CBF ratio of frontal lobe / occipital lobe between patients and

165 controls, where we found no significant difference between the groups (p=0.10). In addition, we

166 compared the ratio of the spatial coefficients of variation of the frontal lobe/occipital lobe for patients

and controls. Again, we found no significant difference between the groups (p=0.091). To demonstrate

the range of values, the mean CBF of the frontal lobe for each individual in our cohort is shown in Figure

169

1.

170

171 Perfusion of other brain regions

172 We assessed the CBF of several brain regions (Frontal lobe, Occipital lobe, Parietal lobe, Temporal lobe,

173 Insula, Cerebellum, Caudate, Putamen, and Thalamus) in trigonocephaly patients as compared to

- 174 controls, using anatomical structural atlas. We observed no significant difference in trigonocephaly
- patients as compared to controls for these regions. The CBF values are summarized in supplemental
- 176 **Table 4.**
- 177

178 Discussion

179 The aim of this study was to investigate CBF in the frontal lobe of patients with trigonocephaly as 180 compared to age-matched controls. Our study suggests that there is no significant difference in CBF in 181 trigonocephaly patients as compared to healthy controls. This is consistent with our research hypothesis 182 that there are no abnormalities in frontal lobe perfusion in trigonocephaly patients aged from 0-18 183 months. This matches the previous findings of a very low risk to develop raised intracranial pressure in 184 the first 18 months of life in trigonocephaly patients and the fact that trigonocephaly patients have normal intracranial volume compared to controls without cerebral pathology. ^{2, 3} 185 186 Currently, patients with a moderate or severe phenotype of trigonocephaly undergo surgery with the 187 aim to reduce restriction of the brain that might cause raised intracranial pressure and hereby improve 188 brain and neurocognitive development as well as to improve aesthetic outcome. 189 However, the functional indication and the efficacy of surgical correction of trigonocephaly has been 190 under debate among craniofacial surgeons since a few years. It is unclear if premature closure of the 191 metopic suture restricts brain development mechanically, especially of the frontal lobe, and whether 192 craniofacial surgery has a positive effect. The neurocognitive findings reported in older children with treated trigonocephaly might represent an intrinsic brain disorder which is not improved with surgery.²¹ 193 194 Our current finding of equal CBF in trigonocephaly patients before surgery and that of control patients 195 suggests the metopic synostosis does not impair CBF in the forebrain below the age of 18 months, and 196 thus supports the hypothesis of intrinsic brain problems rather than that of mechanical restriction of 197 brain development.

In line with the theory that trigonocephaly is mainly an inborn brain disorder are recent studies which
 have shown that some genetic mutations found in patients with trigonocephaly overlap with patients
 with developmental delay disorders. ⁶ Further studies into the microstructure of the brain, using for

example diffusion tension imaging are required to further clarify brain development in trigonocephaly
 patients and understand the underlying pathophysiology.

The previous imaging SPECT studies in trigonocephaly patients had different findings.^{9, 10} This can be explained by different methodology, we used a quantitative methods instead of the qualitative comparison. Also, the studies have reported some left-right asymmetry and none of that was visible in our subjects despite that the ASL signal is usually very stable across the hemispheres.²² At last, Shimoji et al had a different age range (1-9 year).⁹

208 As the largest ASL study of healthy children to date, Carsin-Vu et al showed a mean perfusion of 54.6 209 mL/100gr/min and of 68.4 mL/100gr/min in the frontal lobe of 6 -11 months (n=4) and 12-23 months (n=14) old healthy children respectively, which is roughly similar to our results. ²³ The differences are 210 211 likely to stem from differences in the methodological issues of brain region definition (e.g. manual or 212 automatic ROIs), different ASL sequences types (pulsed, pseudocontinuous), and different ASL 213 acquisition parameters. As the range of our cohort in figure 1 is shown, we expect that age will not have 214 had a significant effect on our results. Due to our limited sample size we did not conduct additional 215 analyses on age and its correlation with cerebral perfusion.

216

217 In this study segmentation was difficult which had an influence on the registration to the MNI atlases to

assess CBF regionally. But we have used the registration of CBF to pseudoCBF that was shown by

219 Mutsaerts in 2018.¹⁷ On top of that, we managed to additionally use a low-degree-of-freedom non-

220 linear component to improve the registration for the deformed skulls thus reaching a better registration.

221 This was evaluated both visually and quantitatively, which is subject of another manuscript.

223 Our study has several limitations. First, our study focusses on a limited number of patients. Therefore, 224 establishing that there is no difference in CBF between trigonocephaly patients and controls with 225 certainty remains difficult. We demonstrated that the range of CBF in both trigonocephaly patients and 226 controls is similar to the range found in previous studies. In our cohort there was no significant 227 difference between mean CBF of the frontal lobe in trigonocephaly patients as compared to controls. 228 We tested these results by both checking the CBF ratio of frontal lobe/occipital lobe in patients as 229 compared to controls and by establishing the ratio of spatial coefficients of variation of the frontal 230 lobe/occipital lobe in patients vs controls, where both tests showed no significant difference. Still, large 231 cohort studies and therefore standardized CBF values in pediatric patients are missing. ASL studies 232 focusing on clinical relevance of differences in CBF range are required for the pediatric setting. 233 234 Second, we were unable to obtain exact-aged matched healthy controls, due to ethical constraints on 235 subjecting healthy children to the anesthesia required for undergoing MRI examination, solely for 236 research purposes. Our control group therefore consisted of patients who underwent MRI examination 237 for clinical reasons, where MRI and clinical course showed no cerebral pathology. Patients with 238 trigonocephaly tend to go younger to MRI because of the visibility disorder which could develop later in 239 life. This age difference might have produced an additional mean CBF difference of -0.3, 0.12, -0.06 240 mL/100g/min (for the three presented models) between the groups, which would have then even

reduced the group difference or not change it significantly.²³

Third, we did not differentiate between gray and white matter when evaluating the mean regional CBF due to the absence of T1-weighted image segmentation. Instead, the mean parenchymal CBF was assessed per region. Gray matter perfusion is around 2 times higher than white matter perfusion in pediatric populations and joint evaluation of gray matter and white matter signals can bias the CBF analysis if gray matter and white matter volumes differ significantly between groups.²³⁻²⁵ However, such

247	volumetric difference between GM and WM is not expected, so we estimate that this had no influence.
248	More advanced analysis that uses T2w and DTI images to aid the segmentation and study the partial
249	volume corrected gray matter CBF is however planned. ²⁶ As the study of Carsin-Vu et al among pediatric
250	patients found no difference in CBF between patients of a different sex or the use of different types of
251	anesthesia, we did not take these factors into account in our analysis. ²³
252	
253	Conclusions
254	In conclusion, our finding of a normal CBF in untreated trigonocephaly patients under the age of 18
255	months as compared to controls supports a more conservative approach to prevent potential
256	overtreatment of patients with trigonocephaly. In addition to the previously reported very low
257	prevalence of papilledema or impaired skull growth, this finding further supports our hypothesis that
258	craniofacial surgery for trigonocephaly is rarely indicated for signs of raised intracranial pressure.
259	

260 Acknowledgements: none

261 **Disclosure**: none

262 Figure Legends

263 References

1. Cornelissen M, den Ottelander B, Rizopoulos D, et al. Increase of prevalence of craniosynostosis.

265 Journal of Cranio-Maxillofacial Surgery 2016;44:1273-1279

- 266 2. Cornelissen MJ, Loudon SE, van Doorn FE, et al. Very Low Prevalence of Intracranial
- 267 Hypertension in Trigonocephaly. *Plast Reconstr Surg* 2017;139:97e-104e
- 268 3. Maltese G, Tarnow P, Wikberg E, et al. Intracranial volume before and after surgical treatment
- 269 for isolated metopic synostosis. *J Craniofac Surg* 2014;25:262-266
- 4. van der Vlugt JJ, van der Meulen JJ, Creemers HE, et al. Cognitive and behavioral functioning in
- 271 82 patients with trigonocephaly. *Plast Reconstr Surg* 2012;130:885-893
- 5. Kelleher MO, Murray DJ, McGillivary A, et al. Behavioral, developmental, and educational
- problems in children with nonsyndromic trigonocephaly. J Neurosurg 2006;105:382-384
- 274 6. Mocquard C, Aillet S, Riffaud L. Recent advances in trigonocephaly. *Neurochirurgie* 2019;65:246-
- 275 251
- 276 7. Grandhi R, Peitz GW, Foley LM, et al. The influence of suturectomy on age-related changes in
- 277 cerebral blood flow in rabbits with familial bicoronal suture craniosynostosis: A quantitative analysis.
- 278 PLoS One 2018;13:e0197296
- 279 8. Doerga PN, Lequin MH, Dremmen MHG, et al. Cerebral blood flow in children with syndromic
- 280 craniosynostosis: cohort arterial spin labeling studies. J Neurosurg Pediatr 2019:1-11
- 281 9. Shimoji T, Shimabukuro S, Sugama S, et al. Mild trigonocephaly with clinical symptoms: analysis
- of surgical results in 65 patients. Childs Nerv Syst 2002;18:215-224
- 10. Sen A, Dougal P, Padhy AK, et al. Technetium-99m-HMPAO SPECT cerebral blood flow study in
- 284 children with craniosynostosis. J Nucl Med 1995;36:394-398

- 285 11. Alsop DC, Detre JA, Golay X, et al. Recommended implementation of arterial spin-labeled
- 286 perfusion MRI for clinical applications: A consensus of the ISMRM perfusion study group and the
- 287 European consortium for ASL in dementia. Magn Reson Med 2015;73:102-116
- 288 12. Yerys BE, Herrington JD, Bartley GK, et al. Arterial spin labeling provides a reliable
- 289 neurobiological marker of autism spectrum disorder. J Neurodev Disord 2018;10:32
- 290 13. Hales PW, d'Arco F, Cooper J, et al. Arterial spin labelling and diffusion-weighted imaging in
- 291 paediatric brain tumours. *Neuroimage Clin* 2019;22:101696
- 292 14. Birgfeld CB, Saltzman BS, Hing AV, et al. Making the diagnosis: metopic ridge versus metopic
- craniosynostosis. J Craniofac Surg 2013;24:178-185
- 15. Mutsaerts H, Petr J, Groot P, et al. ExploreASL: An image processing pipeline for multi-center ASL
- 295 perfusion MRI studies. *Neuroimage* 2020;219:117031
- 296 16. Shi F, Yap PT, Wu G, et al. Infant brain atlases from neonates to 1- and 2-year-olds. *PLoS One*

297 2011;6:e18746

- 298 17. Mutsaerts H, Petr J, Thomas DL, et al. Comparison of arterial spin labeling registration strategies
- 299 in the multi-center GENetic frontotemporal dementia initiative (GENFI). J Magn Reson Imaging
- 300 2018;47:131-140
- 301 18. Ashburner J, Friston KJ. Nonlinear spatial normalization using basis functions. *Hum Brain Mapp* 302 1999;7:254-266
- 303 19. Hammers A, Allom R, Koepp MJ, et al. Three-dimensional maximum probability atlas of the
- human brain, with particular reference to the temporal lobe. *Hum Brain Mapp* 2003;19:224-247
- 20. Team RC. R: A language and environment for statistical computing. R Foundation for Statistical
- 306 Computing. 1.1.442 ed. Vienna, Austria; 2013
- 307 21. van der Meulen J. Metopic synostosis. Childs Nerv Syst 2012;28:1359-1367

- 308 22. Arnaud-Lopez L, Fragoso R, Mantilla-Capacho J, et al. Crouzon with acanthosis nigricans. Further
- delineation of the syndrome. *Clin Genet* 2007;72:405-410
- 310 23. Carsin-Vu A, Corouge I, Commowick O, et al. Measurement of pediatric regional cerebral blood
- flow from 6 months to 15 years of age in a clinical population. *Eur J Radiol* 2018;101:38-44
- 312 24. Kim HG, Lee JH, Choi JW, et al. Multidelay Arterial Spin-Labeling MRI in Neonates and Infants:
- 313 Cerebral Perfusion Changes during Brain Maturation. *AJNR Am J Neuroradiol* 2018;39:1912-1918
- 25. Tortora D, Mattei PA, Navarra R, et al. Prematurity and brain perfusion: Arterial spin labeling
- 315 MRI. Neuroimage Clin 2017;15:401-407
- 26. Feng L, Li H, Oishi K, et al. Age-specific gray and white matter DTI atlas for human brain at 33, 36
- and 39 postmenstrual weeks. *Neuroimage* 2019;185:685-698

320 Tables

321

322 Table 1. Patient Characteristics

	Trigonocephaly	Controls
n	36	16
f:m	11:25	10:06
median	0.50	0.83
IQR	0.30	0.56

324	Table 2. Mixed model on the	perfusion of the frontal	lobe using the structural atlas	(ml/100gr/min)
-----	-----------------------------	--------------------------	---------------------------------	----------------

	Trigonocephaly								
	mean	SE	lower.ci	upper.ci	mean	SE	lower.ci	upper.ci	p-value
Frontal lobe	73.0	2.97	67.0	78.9	70.5	4.45	61.6	79.4	0.6479

Figure 1. Mean CBF (ml/100gr/min) of the frontal lobe in trigonocephaly patients and controls over time





Table 3. Mixed model on the three levels of the frontal lobe using the Hammers Atlas (ml/100gr/min)

	Trigonocephaly				Control				
	mean	mean SE lower.ci upper.ci			mean	SE	lower.ci	upper.ci	
Frontal lobe superior	70.9	2.90	65.1	76.7	68.2	4.34	59.5	76.9	
Frontal lobe middle	68.4	3.09	62.2	74.6	70.1	4.63	60.8	79.4	
Frontal lobe inferior	76.8	2.9	70.9	82.6	75.9	4.5	66.8	84.9	

Supplemental table 4. Perfusion per brain region from the structural atlas in ml/100gr/min

	Trigonocephaly					Control				
Brain Region	mean	sd	se	lower.ci	upper.ci	mean	sd	se	lower.ci	upper.ci
Frontal	72.97	16.77	1.98	69.03	76.91	70.51	19.97	3.53	63.31	77.71
Occipital	80.32	20.95	2.47	75.40	85.25	71.60	18.20	3.22	65.04	78.16
Parietal	77.49	18.37	2.16	73.18	81.81	72.94	17.35	3.07	66.68	79.20
Temporal	71.90	15.91	1.88	68.16	75.64	69.93	19.87	3.51	62.77	77.10
Insula	79.93	15.96	1.88	76.18	83.68	73.34	21.95	3.88	65.43	81.25
Cerebellum	73.96	15.67	1.85	70.28	77.64	66.47	21.88	3.87	58.58	74.36
Caudate	63.25	14.04	1.65	59.95	66.55	56.20	15.61	2.76	50.57	61.83
Putamen	78.93	15.20	1.79	75.36	82.50	70.88	25.23	4.46	61.79	79.98
Thalamus	93.24	25.25	2.98	87.30	99.17	80.03	28.15	4.98	69.88	90.18
	•									