




# **Testing CVC and CKM unitarity via superallowed nuclear $\beta$ decay**

**J.C. Hardy and I.S. Towner**

**Cyclotron Institute  
Texas A&M University**



**H.I. Park  
V.E. Jacob  
L. Chen  
V. Horvat  
N. Nica  
M. Bencomo**

# SUPERALLOWED $0^+ \rightarrow 0^+$ BETA DECAY

## BASIC WEAK-DECAY EQUATION


$$ft = \frac{K}{G_V^2 <\tau>^2}$$

$f$  = statistical rate function:  $f(Z, Q_{EC})$

$t$  = partial half-life:  $t_{1/2}$

$G_V$  = vector coupling constant

$<\tau>$  = Fermi matrix element



# SUPERALLOWED $0^+ \rightarrow 0^+$ BETA DECAY

## BASIC WEAK-DECAY EQUATION


$$ft = \frac{K}{G_V^2 <\tau>^2}$$

$f$  = statistical rate function:  $f(Z, Q_{EC})$

$t$  = partial half-life:  $t_{1/2}$

$G_V$  = vector coupling constant

$<\tau>$  = Fermi matrix element



## INCLUDING RADIATIVE AND ISOSPIN-SYMMETRY-BREAKING CORRECTIONS

$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2 (1 + \Delta_R)}$$

# SUPERALLOWED $0^+ \rightarrow 0^+$ BETA DECAY

## BASIC WEAK-DECAY EQUATION


$$ft = \frac{K}{G_V^2 \langle \tau \rangle^2}$$

$f$  = statistical rate function:  $f(Z, Q_{EC})$

$t$  = partial half-life:  $f(t_{1/2}, BR)$

$G_V$  = vector coupling constant

$\langle \tau \rangle$  = Fermi matrix element



## INCLUDING RADIATIVE AND ISOSPIN-SYMMETRY-BREAKING CORRECTIONS

$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2 (1 + \Delta_R)}$$

$f(Z, Q_{EC})$

~1.5%

$f(\text{nuclear structure})$

0.3-1.5%

$f(\text{interaction})$

~2.4%

# SUPERALLOWED $0^+ \rightarrow 0^+$ BETA DECAY

## BASIC WEAK-DECAY EQUATION


$$ft = \frac{K}{G_v^2 <\tau>^2}$$

$f$  = statistical rate function:  $f(Z, Q_{EC})$

$t$  = partial half-life:  $f(t_{1/2}, BR)$

$G_v$  = vector coupling constant

$<\tau>$  = Fermi matrix element



## INCLUDING RADIATIVE AND ISOSPIN-SYMMETRY-BREAKING CORRECTIONS

$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_v^2 (1 + \Delta_R)}$$

$f(Z, Q_{EC})$

~1.5%

$f(\text{nuclear structure})$

0.3-1.5%

$f(\text{interaction})$

~2.4%

## THEORETICAL UNCERTAINTIES

0.05 – 0.10%

# WHAT CAN WE LEARN?

## WHAT CAN WE LEARN?

FROM A SINGLE TRANSITION

Experimentally  
determine  $G_v^2(1 + \frac{1}{R})$

$$\mathcal{F}t = ft(1 + \frac{1}{R})[1 - (\frac{c}{c_{NS}} - \frac{1}{R})] = \frac{K}{2G_v^2(1 + \frac{1}{R})}$$

# WHAT CAN WE LEARN?

## FROM A SINGLE TRANSITION

Experimentally  
determine  $G_v^2(1 + \frac{r}{R})$

$$\mathcal{F}t = ft(1 + \frac{r}{R})[1 - (\frac{c}{c} - \frac{ns}{ns})] = \frac{K}{2G_v^2(1 + \frac{r}{R})}$$

## FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

Validate the correction  
terms

Test for presence of  
a Scalar current

$\mathcal{F}t$  values constant

# WHAT CAN WE LEARN?

## FROM A SINGLE TRANSITION

Experimentally  
determine  $G_v^2(1 + \frac{K}{R})$

$$\mathcal{F}t = ft(1 + \frac{K}{R})[1 - (\frac{c}{c} - \frac{ns}{ns})] = \frac{K}{2G_v^2(1 + \frac{K}{R})}$$

## FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

Validate the correction  
terms

Test for presence of  
a Scalar current

$\mathcal{F}t$  values constant

## WITH CVC VERIFIED

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

weak eigenstates      Cabibbo Kobayashi Maskawa (CKM) matrix      mass eigenstates

Obtain precise value of  $G_v^2(1 + \frac{K}{R})$   
Determine  $V_{ud}^2$

$$V_{ud}^2 = G_v^2/G^2$$

# WHAT CAN WE LEARN?

## FROM A SINGLE TRANSITION

Experimentally determine  $G_v^2(1 + \frac{K}{R})$

$$\mathcal{F}t = ft(1 + \frac{K}{R})[1 - (\frac{c}{c} - \frac{ns}{ns})] = \frac{K}{2G_v^2(1 + \frac{K}{R})}$$

## FROM MANY TRANSITIONS

Test Conservation of the Vector current (CVC)

Validate the correction terms

Test for presence of a Scalar current

$\mathcal{F}t$  values constant

## WITH CVC VERIFIED

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

weak eigenstates      Cabibbo Kobayashi Maskawa (CKM) matrix      mass eigenstates

Obtain precise value of  $G_v^2(1 + \frac{K}{R})$   
Determine  $V_{ud}^2$

Test CKM unitarity

$$V_{ud}^2 = G_v^2/G^2$$

$$V_{ud}^2 + V_{us}^2 + V_{ub}^2 = 1$$

# WHAT CAN WE LEARN?

FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \frac{R}{R})$

$$\mathcal{F}t = ft(1 + \frac{R}{R})[1 - (\frac{c}{c} - \frac{ns}{ns})] = \frac{K}{2G_V^2(1 + \frac{R}{R})}$$

FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

Validate the correction  
terms

Test for presence of  
a Scalar current

$\mathcal{F}t$  values constant

WITH CVC VERIFIED

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$


weak eigenstates      Cabibbo Kobayashi Maskawa (CKM) matrix      mass eigenstates

Obtain precise  
Determine  
**ONLY POSSIBLE IF PRIOR  
CONDITIONS SATISFIED**

$$V_{ud}^2 = G_V^2/G^2$$

$$V_{ud}^2 + V_{us}^2 + V_{ub}^2 = 1$$



# WORLD DATA FOR $0^+ \rightarrow 0^+$ DECAY, 2014



- 8 cases with  $ft$ -values measured to <0.05% precision; 6 more cases with 0.05-0.3% precision.
- ~220 individual measurements with compatible precision

$$ft = f't(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$


# WORLD DATA FOR $0^+ \rightarrow 0^+$ DECAY, 2014




- 8 cases with  $ft$ -values measured to <0.05% precision; 6 more cases with 0.05-0.3% precision.
- ~220 individual measurements with compatible precision

$$\mathcal{F}t = \textcolor{red}{ft} (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2 (1 + \Delta_R)}$$

# WORLD DATA FOR $0^+ \rightarrow 0^+$ DECAY, 2014




- 8 cases with  $ft$ -values measured to <0.05% precision; 6 more cases with 0.05-0.3% precision.
- ~220 individual measurements with compatible precision




$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2 (1 + \Delta_R)}$$

# WORLD DATA FOR $0^+ \rightarrow 0^+$ DECAY, 2014



- 8 cases with  $ft$ -values measured to <0.05% precision; 6 more cases with 0.05-0.3% precision.
- ~220 individual measurements with compatible precision

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2 (1 + \Delta_R)}$$




# TESTS OF $\delta_c$ CALCULATIONS

## A. Agreement with CVC:

$\mathcal{F}_t$  values have been calculated with different models for  $\delta_c$ , then tested for consistency. Normalized  $\chi^2$  and confidence levels are shown.

Model	$\chi^2/N$	CL(%)
SM-SW	1.37	17





# TESTS OF $\delta_c$ CALCULATIONS

## A. Agreement with CVC:

$\mathcal{F}_t$  values have been calculated with different models for  $\delta_c$ , then tested for consistency. Normalized  $\chi^2$  and confidence levels are shown.

Model	$\chi^2/N$	CL(%)
SM-SW	1.37	17
SM-HF	6.38	0






# TESTS OF $\delta_c$ CALCULATIONS

## A. Agreement with CVC:

$\mathcal{F}_t$  values have been calculated with different models for  $\delta_c$ , then tested for consistency. Normalized  $\chi^2$  and confidence levels are shown.

Model	$\chi^2/N$	CL(%)
SM-SW	1.37	17
SM-HF	6.38	0
DFT	4.26	0






# TESTS OF $\delta_c$ CALCULATIONS

## A. Agreement with CVC:


$\mathcal{F}_t$  values have been calculated with different models for  $\delta_c$ , then tested for consistency. Normalized  $\chi^2$  and confidence levels are shown.

Model	$\chi^2/N$	CL(%)
SM-SW	1.37	17
SM-HF	6.38	0
DFT	4.26	0
RHF-RPA	4.91	0
RH-RPA	3.68	0



# TESTS OF $\delta_c$ CALCULATIONS


## B. Measurements of mirror superallowed transitions:



# TESTS OF $\delta_c$ CALCULATIONS

## B. Measurements of mirror superallowed transitions:

$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})]$$




$$\begin{aligned} \frac{ft_A}{ft_B} &= \frac{(1 + \delta'^B_R)[1 - (\delta^B_c - \delta^B_{NS})]}{(1 + \delta'^A_R)[1 - (\delta^A_c - \delta^A_{NS})]} \\ &= 1 + (\delta'^B_R - \delta'^A_R) + (\delta^B_{NS} - \delta^A_{NS}) - (\delta^B_c - \delta^A_c) \end{aligned}$$


# TESTS OF $\delta_c$ CALCULATIONS

## B. Measurements of mirror superallowed transitions:


$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})]$$




$$\begin{aligned} \frac{ft_A}{ft_B} &= \frac{(1 + \delta'^B_R)[1 - (\delta^B_c - \delta^B_{NS})]}{(1 + \delta'^A_R)[1 - (\delta^A_c - \delta^A_{NS})]} \\ &= 1 + (\delta'^B_R - \delta'^A_R) + (\delta^B_{NS} - \delta^A_{NS}) - (\delta^B_c - \delta^A_c) \end{aligned}$$





# ACCESSIBLE MIRROR PAIRS OF SUPERALLOWED DECAYS





# ACCESSIBLE MIRROR PAIRS OF SUPERALLOWED DECAYS





# PRECISION DECAY MEASUREMENTS AT TAMU






# PRECISION DECAY MEASUREMENTS AT TAMU






# PRECISION DECAY MEASUREMENTS AT TAMU




# BETA-DECAY BRANCHING OF $^{38}\text{Ca}$





# BETA-DECAY BRANCHING OF $^{38}\text{Ca}$




# BETA-DECAY BRANCHING OF $^{38}\text{Ca}$



$$\frac{N_{\gamma_1\beta}}{N_\beta} = \frac{N_o R_1 \epsilon_{\beta_1} \epsilon_{\gamma_1}}{N_o \epsilon_{\beta_{\text{tot}}}}$$





# BETA-DECAY BRANCHING OF $^{38}\text{Ca}$




$$\frac{N_{\gamma_1\beta}}{N_\beta} = \frac{N_o R_1 \epsilon_{\beta_1} \epsilon_{\gamma_1}}{N_o \epsilon_{\beta_{\text{tot}}}}$$

$$R_1 = \frac{N_{\gamma_1\beta}}{N_\beta \epsilon_{\gamma_1} \epsilon_{\beta_1}} k$$





# BETA-DECAY BRANCHING OF $^{38}\text{Ca}$






$$\frac{N_{\gamma_1\beta}}{N_\beta} = \frac{N_o R_1 \epsilon_{\beta_1} \epsilon_{\gamma_1}}{N_o \epsilon_{\beta_{\text{tot}}}}$$

$$R_1 = \frac{N_{\gamma_1\beta}}{N_\beta \epsilon_{\gamma_1} \epsilon_{\beta_1}} k$$

$k = f(\text{dead-time, pile-up, coincidence summing}) = 1.0391(23)$



# BETA-DECAY BRANCHING OF $^{38}\text{Ca}$



$$\frac{N_{\gamma_1\beta}}{N_\beta} = \frac{N_o R_1 \epsilon_{\beta_1} \epsilon_{\gamma_1}}{N_o \epsilon_{\beta_{\text{tot}}}}$$


$$R_1 = \frac{N_{\gamma_1\beta}}{N_\beta \epsilon_{\gamma_1} \epsilon_{\beta_1}} k$$

$k = f(\text{dead-time, pile-up, coincidence summing}) = 1.0391(23)$


# TESTS OF $\delta_c$ CALCULATIONS

## B. Measurements of mirror superallowed transitions:

$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})]$$




$$\begin{aligned} \frac{ft_A}{ft_B} &= \frac{(1 + \delta'^B_R)[1 - (\delta^B_c - \delta^B_{NS})]}{(1 + \delta'^A_R)[1 - (\delta^A_c - \delta^A_{NS})]} \\ &= 1 + (\delta'^B_R - \delta'^A_R) + (\delta^B_{NS} - \delta^A_{NS}) - (\delta^B_c - \delta^A_c) \end{aligned}$$




# TESTS OF $\delta_c$ CALCULATIONS

## B. Measurements of mirror superallowed transitions:

$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})]$$



$$\begin{aligned} \frac{ft_A}{ft_B} &= \frac{(1 + \delta'^B_R)[1 - (\delta^B_c - \delta^B_{NS})]}{(1 + \delta'^A_R)[1 - (\delta^A_c - \delta^A_{NS})]} \\ &= 1 + (\delta'^B_R - \delta'^A_R) + (\delta^B_{NS} - \delta^A_{NS}) - (\delta^B_c - \delta^A_c) \end{aligned}$$



## RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \Delta_R)$

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

FROM A SINGLE TRANSITION


Experimentally  
determine  $G_V^2(1 + \Delta_R)$

$$\bar{\tau}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

$G_V$  constant to  $\pm 0.013\%$



$$\begin{aligned}\bar{\tau}t &= 3072.3(7) \\ G_V(1+\Delta_R)^{1/2}/(hc)^3 &= 1.14958(15) \\ &\times 10^{-5} \text{ GeV}^{-2}\end{aligned}$$

$$\chi^2/\nu = 0.6$$

# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

## FROM A SINGLE TRANSITION

Experimentally  
determine  $G_v^2(1 + \Delta_R)$

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_v^2(1 + \Delta_R)}$$

## FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

Validate correction terms

$G_v$  constant to  $\pm 0.013\%$

# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \Delta_R)$


$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

Validate correction terms

$G_V$  constant to  $\pm 0.013\%$



# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \Delta_R)$


$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2 (1 + \Delta_R)}$$

FROM MANY TRANSITIONS


Test Conservation of  
the Vector current (CVC)

Validate correction terms ✓

Model	$\chi^2/N$	CL(%)
SM-SW	1.37	17
SM-HF	6.38	0
DFT	4.26	0
RHF-RPA	4.91	0
RH-RPA	3.68	0



$G_V$  constant to  $\pm 0.013\%$



# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

## FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \Delta_R)$

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

## FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

Validate correction terms ✓

Test for Scalar current

$G_V$  constant to  $\pm 0.013\%$

# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

## FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \Delta_R)$

$$\mathcal{F}t = ft (1 + \delta'_R) [1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2 (1 + \Delta_R)}$$

## FROM MANY TRANSITIONS


Test Conservation of  
the Vector current (CVC)

Validate correction terms ✓

Test for Scalar current

$G_V$  constant to  $\pm 0.013\%$

limit,  $C_s/C_v = 0.0014 (13)$



# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

## FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \Delta_R)$

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

## FROM MANY TRANSITIONS



Test Conservation of  
the Vector current (CVC)

Validate correction terms ✓

Test for Scalar current

$G_V$  constant to  $\pm 0.013\%$

limit,  $C_s/C_v = 0.0014 (13)$



# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

## FROM A SINGLE TRANSITION

Experimentally  
determine  $G_V^2(1 + \Delta_R)$

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

## FROM MANY TRANSITIONS

Test Conservation of  
the Vector current (CVC)

Validate correction terms ✓

Test for Scalar current

$G_V$  constant to  $\pm 0.013\%$

limit,  $C_s/C_v = 0.0014 (13)$

## WITH CVC VERIFIED

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

weak eigenstates      mass eigenstates

Obtain precise value of  $G_V^2(1 + \Delta_R)$   
Determine  $V_{ud}^2$

$$V_{ud}^2 = G_V^2/G_\mu^2 = 0.94900 \pm 0.00042$$

Cabibbo-Kobayashi-Maskawa matrix

# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

## FROM A SINGLE TRANSITION

Experimentally determine  $G_V^2(1 + \Delta_R)$

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

## FROM MANY TRANSITIONS

Test Conservation of the Vector current (CVC)

Validate correction terms ✓

Test for Scalar current

$G_V$  constant to  $\pm 0.013\%$

limit,  $C_s/C_v = 0.0014$  (13)

## WITH CVC VERIFIED


$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

weak eigenstates      mass eigenstates

Cabibbo-Kobayashi-Maskawa matrix

Obtain precise value of  $G_V^2(1 + \Delta_R)$   
Determine  $V_{ud}^2$

$$V_{ud}^2 = G_V^2/G_\mu^2 = 0.94900 \pm 0.00042$$



# RESULTS FROM $0^+ \rightarrow 0^+$ DECAY

## FROM A SINGLE TRANSITION

Experimentally determine  $G_V^2(1 + \Delta_R)$

$$\mathcal{F}t = ft(1 + \delta'_R)[1 - (\delta_c - \delta_{NS})] = \frac{K}{2G_V^2(1 + \Delta_R)}$$

## FROM MANY TRANSITIONS

Test Conservation of the Vector current (CVC)

Validate correction terms ✓

Test for Scalar current

$G_V$  constant to  $\pm 0.013\%$

limit,  $C_s/C_v = 0.0014$  (13)

## WITH CVC VERIFIED

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

weak eigenstates      mass eigenstates  
Cabibbo-Kobayashi-Maskawa matrix

Obtain precise value of  $G_V^2(1 + \Delta_R)$

Determine  $V_{ud}^2$

$$V_{ud}^2 = G_V^2/G_\mu^2 = 0.94900 \pm 0.00042$$

Test CKM unitarity

$$V_{ud}^2 + V_{us}^2 + V_{ub}^2 = 0.99992 \pm 0.00048$$

# CURRENT STATUS OF CKM UNITARITY

$$V_{ud}^2 + V_{us}^2 + V_{ub}^2 = 0.99992 \pm 0.00048$$

$V_{ud}^2$  nuclear decays  
muon decay  
 $0.94900 \pm 0.00042$

$V_{us}^2$  kaon decays  
 $0.05090 \pm 0.00022$

$V_{ub}^2$  B decays  
 $0.00002 \pm 0.00001$



# CURRENT STATUS OF CKM UNITARITY

$$V_{ud}^2 + V_{us}^2 + V_{ub}^2 = 0.99992 \pm 0.00048$$

$V_{ud}^2$  nuclear decays  
 $V_{ud}^2$  muon decay  
 $0.94900 \pm 0.00042$

$V_{us}^2$  kaon decays  
 $0.05090 \pm 0.00022$

$V_{ub}^2$  B decays  
 $0.00002 \pm 0.00001$



## FINAL REMARK ON $V_{us}$

Kaon decay yields two independent determinations of  $V_{us}$ :

- 1) Semi-leptonic  $K \rightarrow \pi \ell \nu_\ell$  decay ( $K_{\ell 3}$ ) yields  $|V_{us}|$ .
- 2) Pure leptonic decays  $K^+ \rightarrow \mu^+ \nu_\mu$  and  $\pi^+ \rightarrow \mu^+ \nu_\mu$  together yield  $|V_{us}| / |V_{ud}|$ .

Both require lattice calculations of form factors to obtain their result.

Until March 2014 these gave highly consistent results for  $|V_{us}|$ .

## FINAL REMARK ON $V_{us}$

Kaon decay yields two independent determinations of  $V_{us}$ :

- 1) Semi-leptonic  $K \rightarrow \pi \ell \nu_\ell$  decay ( $K_{\ell 3}$ ) yields  $|V_{us}|$ .
- 2) Pure leptonic decays  $K^+ \rightarrow \mu^+ \nu_\mu$  and  $\pi^+ \rightarrow \mu^+ \nu_\mu$  together yield  $|V_{us}| / |V_{ud}|$ .

Both require lattice calculations of form factors to obtain their result.

Until March 2014 these gave highly consistent results for  $|V_{us}|$ .

**BUT**, Bazavov et al. [PRL 112, 112001 (2014)] produced a new lattice calculation of the form factor used for  $K_{\ell 3}$  decays.

Their new result for  $|V_{us}|$  is inconsistent with the  $|V_{us}| / |V_{ud}|$  result and, when combined with the superallowed result for  $|V_{ud}|$ , leads to a unitarity sum over two standard deviations below 1.

Stay tuned ...

## SUMMARY AND OUTLOOK

1. Analysis of superallowed  $0^+ \rightarrow 0^+$  nuclear  $\beta$  decay is shown to confirm CVC and thus yield  $V_{ud} = 0.97417(21)$ .
2. The three other experimental methods for determining  $V_{ud}$  yield consistent results, but are less precise by a factor of 8 or more.
3. The current value for  $V_{ud}$ , when combined with  $V_{us}$  and  $V_{ub}$ , satisfies CKM unitarity to 0.05%.

## SUMMARY AND OUTLOOK

1. Analysis of superallowed  $0^+ \rightarrow 0^+$  nuclear  $\beta$  decay is shown to confirm CVC and thus yield  $V_{ud} = 0.97417(21)$ .
2. The three other experimental methods for determining  $V_{ud}$  yield consistent results, but are less precise by a factor of 8 or more.
3. The current value for  $V_{ud}$ , when combined with  $V_{us}$  and  $V_{ub}$ , satisfies CKM unitarity to 0.05%.

4. The largest contribution to the  $V_{ud}$  uncertainty is from the inner radiative correction. Isospin symmetry-breaking corrections in nuclei are the second largest.
5. These symmetry-breaking corrections have been tested by requiring consistency among 14 known transitions (CVC) and by comparing them with mirror transition pairs. One set of corrections passes both tests.
6. Further tests on mirror pairs are possible and are planned. This requires precise branching-ratio measurements of  $T_z = -1$  parent decays.